



SYNAGIS WEB BASED FORM

Submit online or fax completed form to: 855-207-0250

For questions regarding this prior authorization, call 866-773-0695

Prior Authorization Vendor for ND Medicaid

Note:

- Synagis season will be October 19th through April 21st
- Providers will choose when to start dosing Synagis based on prevalence of RSV in the community
- Clinicians may administer up to a maximum of 5 monthly doses during the RSV season.
- Qualifying infants born during the RSV season may require fewer doses.

TO BE COMPLETED BY PRESCRIBER

Recipient Medicaid ID Number	Recipient Date of Birth	Prescriber NPI	Prescriber Fax Number
Billing Facility NPI	Billing Facility Name		ICD-10 code

Is this request for dose from the current RSV season, last RSV season, or both (please select one)?

Previous Year's RSV season
 Current Year's RSV Season
 Both

Diagnosis (qualification for Synagis)

___ **Prematurity**

<29 weeks, 0 days gestational age – Synagis allowed if younger than 12 months of age at start of RSV season (max of 5 doses)

Gestational Age (e.g. 28 weeks, 4 days)

Weeks _____ Days _____

___ **Chronic Lung Disease of Prematurity (CLD)** – Child ≤12 months old with gestational age <32 weeks, 0 days and requires supplemental oxygen >21% for at least the first 28 days after birth.

___ **Chronic Lung Disease of Prematurity (CLD)** – Child ≤24 months old with gestational age <32 weeks, 0 days and requires supplemental oxygen >21% for at least the first 28 days after birth and continues to receive medical support within six months before the start of RSV season.

___ Supplemental Oxygen

___ Diuretic

___ Chronic corticosteroid therapy

___ **Congenital Heart Disease (CHD)**

Child ≤12 months old with hemodynamically significant cyanotic or acyanotic CHD

Medical Therapy Required _____

*children less than 24 months who undergo cardiac transplantation during RSV season may be considered for prophylaxis.

___ **Neuromuscular disease** (may be considered for prophylaxis during the first year of life)

___ **Pulmonary abnormalities** (may be considered for prophylaxis during the first year of life)

___ **Profoundly Immunocompromised children** (children <24 months of age may be considered for prophylaxis during the RSV season)

*Accessed online at pediatrics.aappublications.org